

<https://helda.helsinki.fi>

Onthophagus pilauco sp. nov. (Coleoptera, Scarabaeidae) :
evidence of beetle extinction in the Pleistocene- Holocene
transition in Chilean Northern Patagonia

Tello, Francisco

2021-06-15

Tello , F , Verdu , J R , Rossini , M & Zunino , M 2021 , ' Onthophagus pilauco sp. nov.
(Coleoptera, Scarabaeidae) : evidence of beetle extinction in the Pleistocene- Holocene
transition in Chilean Northern Patagonia ' , ZooKeys , no. 1043 , pp. 133-145 . <https://doi.org/10.3897/zookeys.1043>.

<http://hdl.handle.net/10138/332675>

<https://doi.org/10.3897/zookeys.1043.61706>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Onthophagus pilauco sp. nov. (Coleoptera, Scarabaeidae): evidence of beetle extinction in the Pleistocene–Holocene transition in Chilean Northern Patagonia

Francisco Tello, José R. Verdú, Michele Rossini, Mario Zunino

1 Transdisciplinary Center for Quaternary Research (TAQUACH), Universidad Austral de Chile, Valdivia, Chile
2 I.U.I. CIBIO, Universidad de Alicante, Alicante, Spain
3 Finnish Museum of Natural History (LUOMUS), University of Helsinki, Pohjoinen Rautatiekatu 13, Helsinki, 00014, Finland
4 Scuola di Biodiversità, Polo universitario Asti Studi Superiori, Asti, Italy

Corresponding author: Francisco Tello (ftello@yahoo.cl)

Academic editor: Andrey Frolov | Received 5 December 2020 | Accepted 18 April 2021 | Published 15 June 2021

<http://zoobank.org/B38C0C9B-D6AA-4473-9839-C6DEF8E9FD3B>

Citation: Tello F, Verdú JR, Rossini M, Zunino M (2021) *Onthophagus pilauco* sp. nov. (Coleoptera, Scarabaeidae): evidence of beetle extinction in the Pleistocene–Holocene transition in Chilean Northern Patagonia. ZooKeys 1043: 133–145. <https://doi.org/10.3897/zookeys.1043.61706>

Abstract

The South American Pleistocene–Holocene transition has been characterized by drastic climatic and diversity changes. These rapid changes induced one of the largest and most recent extinctions in the megafauna at the continental scale. However, examples of the extinction of small animals (e.g., insects) are scarce, and the underlying causes of the extinction have been little studied. In this work, a new extinct dung beetle species is described from a late Pleistocene sequence (~15.2 k cal yr BP) at the paleoarcheological site Pilauco, Chilean Northern Patagonia. Based on morphological characters, this fossil is considered to belong to the genus *Onthophagus* Latreille, 1802 and named *Onthophagus pilauco* sp. nov. We carried out a comprehensive revision of related groups, and we analyzed the possible mechanism of diversification and extinction of this new species. We hypothesized that *Onthophagus pilauco* sp. nov. diversified as a member of the *tesculatus* species-complex following migration processes related to the Great American Biotic Interchange (~3 Ma). The extinction of *O. pilauco* sp. nov. may be related to massive defaunation and climatic changes recorded in the Pleistocene–Holocene transition (12.8 k cal yr BP). This finding is the first record of this genus in Chile, and provides new evidence to support the collateral-extinction hypothesis related to the defaunation.

Keywords

Dung beetle, extinction, fossil beetles, new species, Pleistocene, South America

Materials and methods

Abbreviations

CEMT	Seção de Entomologia da Coleção Zoológica, Universidade Federal de Mato, Grosso, Cuiabá, Brazil;
CMNC	Canadian Museum of Nature, Gatineau, Quebec, Canada;
MPDO	Museo Pleistocénico de Osorno, Osorno, Chile;
MZ	Mario Zunino private collection, Asti, Italy;
MZUF	Museo di Storia Naturale dell'Università di Firenze, Florence, Italy;
NMPC	Národní Muzeum, Prague, Czech Republic.

Drawings and determination of fossil remains

The fossil beetle remains were recovered from the sediment using an adaptation of the water flotation technique described by Hoganson et al. (1989) (see also Tello and Torres 2020). The remains were collected from grid 18AC, at an elevation of 384 cm in bed PB-7. The age span of this bed is ~16.0 to 14.0 k cal yr BP (Pino et al. 2020). The taxonomic placement suggested for the fossil was made after detailed examination and a comparison with multiple modern specimens of South American *Onthophagus* species deposited in the CEMT, CMNC, MZUF, MZ and NMPC collections. For the taxonomic nomenclature, we followed Rossini et al. (2018a, b). Figure 1A was obtained using R software v4.0.3. Figure 2A, 2B was obtained using a scanning electron microscope (variable pressure, EVO MC10), and Fig. 2C, D was obtained using a Leica M205C camera. All figures were processed using Adobe Photoshop 2019 CC. Type material is deposited in the MPDO insect collection, Osorno, Chile.

Results

Systematic paleontology

Order Coleoptera Linnaeus, 1758
 Suborder Polyphaga Emery, 1886
 Family Scarabaeidae Latreille, 1802
 Genus *Onthophagus* Latreille, 1802

Type species *Onthophagus pilauco* sp. nov.

<http://zoobank.org/9B203D54-E27A-432D-8AC8-ED7C33ED2F83>

Fig. 2

Description. **Holotype**. Male, minor form. Clypeus sub-trapezoidal and slightly elongated forward, with anterior margin narrowly and slightly re-entrant, head margin barely




Figure 2. Holotype of *Onthophagus pilauco* nov. sp. A dorsal, B ventral and C frontal views, D detail of the microsculpture.

sinuated at the clypeo-genal junction. Fronto-clypeal region without carina, frons with two close, weak tubercles, strongly advanced in position, in line with the anterior margin of the eyes (Fig. 2A, C). Head surface very finely and evenly punctate. Latero-clypeal region with deeper ocellate punctures. Color dark with metallic green to bronze sheen (Fig. 2D). Pronotum and elytra not found.

Female unknown.

Diagnosis. *Onthophagus pilauco* nov. sp. is considered to be a close relative of *O. confusus* Soucomont, 1932 and *O. insularis* Boheman, 1858, as it shares the following morphological characters with these species: sub-trapezoidal shape of the clypeus; small, slightly deeper cephalic punctation, coupled with very shallow wrinkles in proximity to the genal and clypeal margins. Although the fronto-clypeal region is significantly damaged, there is no indication of a possible carina.

Proposed English and Spanish vernacular names. Pilauco dung beetle (EN) and estercolero de Pilauco (ES).

Etymology. The name of the new species refers to the archeopaleontological site from which the fossil remains were collected.

Discussion

From our observations, the fossil remains found at Pilauco correspond to a new and extinct species of the genus *Onthophagus* closely related to the *hircus* group. *Onthophagus pilauco* sp. nov. represents the first record of an endemic species of this genus in Chile. Moreover, this record brings new evidence of beetle extinction related to the Pleistocene–Holocene transition and massive defaunation after a possible cosmic impact and/or YD cooling reversal events.

Morphological delimitation of the fossil record

Despite the beetle remains only being represented by cephalic parts (clypeus, right gena and frons; fronto-clypeal region partly damaged; left gena absent; see Fig. 2A–D) it is clear that they belong to the genus *Onthophagus*. Close scrutiny of the fossil remains, along with an extensive analysis of multiple specimens belonging to extant American *Onthophagus* led us to assign *O. pilauco* to the *O. hircus* group, and more precisely to the *resculatus* species-complex (Rossini et al. 2018a). Cephalic horn-like tubercles may indicate that the remains belong to a male specimen, probably a minor form. The physical location of the cephalic tubercles is rather unique in the modern American *Onthophagus* fauna. They rise in a very advanced position, in line with the anterior margin of the eyes, and are quite close to each other. The combination of these two characteristics has only ever been found in an undescribed *Onthophagus* from Costa Rica, which was included in the same species group, but in a different taxonomic complex (Rossini, pers. comm. 2020). Close cephalic tubercles are also found in female specimens of species belonging to the *Onthophagus* *clypeatus*, *dicranus* and *mirabilis* groups, but they are always situated at the front, and never as advanced as in *O. pilauco*. Also, the shape of the clypeus in these females is always triangular or evenly curved, terminating at the apex with a margin slightly to distinctly emarginated (with two obtuse teeth).

Hypothesis for the speciation and extinction of *Onthophagus pilauco*

Comprehensive knowledge of the Chilean beetle fauna suggests that only a few species (nine) can be considered to be exclusively associated with dung resources (González-Chang and Pinochet 2015). Thus, six species have been assigned to the Scarabaeidae family (excluding the saprophagous Aphodiinae species): two species belonging to the Deltotilini tribe: *Megathopa villosa* Escholtz and *Scybalophagus rugifrons* Blanchard; and four species with uncertain taxonomic position: *Tasasov and Dimitrov* 2016): *Tesserodoniella elguetae* de Mello & Halter, *T. meridionalis* de Mello & Halter, *Homocopris punctatissimus* (Curtis), and *H. torulosus* (Escholtz) (González-Chang and Pinochet 2015). The placement of the genus *Homocopris* as a suprageneric group requires further research. Additionally, three dung beetle species

belong to the family Geotrupidae Latreille, 1802, represented in Chile by the subfamily Taurocerastinae. *Erickius costulatus* Germain, *F. variolosus* Germain and *Taurocerastes patagonicus*, which are distributed across central and southern Chile.

The Deltotrichini group is distributed in the pantropical zone, being especially diversified in the northern areas of South America. The *Tesserodonia* is related to the Australian genera *Tasserodonta* and *Aptenocanthus* and their ancestors probably originated from the Gondwana supercontinent (Vaz-De-Mello and Halter 2006). In contrast, the extant American *Anthrophagus* is a result of migrations to the continent by intercontinental connections, the current remnants of which are known as Beringia components. These migratory processes occurred at different times and under different geographical and climatic conditions, and involved different ancestors, all belonging to the subgenus *Anthrophagus* sensu stricto (Rossini et al. 2018a, b, Halter et al. 2019, Zunino and Halter 2019). Additionally, after the definitive closure of the Isthmus of Panama (~3 Ma), which eliminated natural barriers, Central America became a permanent bridge from one continent to another, improving migratory conditions for large animals and dung beetles. Therefore, we suggest that the ancestors of *O. pilauco* migrated to South America during the Great American Biotic Interchange, following large mammals at the end of the Pliocene (3 Ma). This migratory-mechanism has been suggested for the extant related group, which arrived by crossing the Andes via the Huancabamba Depression, similarly to other extinct dung beetles (e.g., *Phanaeus violaceus*; Zunino, 2013). Intra-continental migratory patterns have also been reported in extant species of dung beetles that have rapidly colonized new habitats following cattle migrations (e.g., *Digitonthophagus gazellae* (Fabricius, 1787); see Noriega et al. 2020). Thus, ancestors of the *costulatus* species-complex diverged in situ in Chilean areas, resulting in the evolution of *O. pilauco* (Fig. 3). This speciation hypothesis is supported by several studies on Pleistocene Patagonian landscapes (~180 and 26 ka) that have suggested that the rapid contraction and expansion of ice cover has induced drastic changes in biotic distributions and prompted diversification in different groups of organisms (e.g., in amphibians: Nuñez et al. 2020; in mammals: Himes et al. 2008). Moreover, the presence of extant endemic species belonging to the *costulatus* species-complex in western Ecuador and northwestern Peru (*O. panofskyi* and *O. insularis*; Rossini et al. 2018a) suggests that *O. pilauco* could be a species endemic to southern Chile.

On the other hand, the extinction of large animals in South American Pleistocene environments has invoked multiple climate- and human-activity-related hypotheses, and interactions between them (Barnosky et al. 2004). The causes of small animal (<60 kg) extinctions, and the implications of changes in species compositions and distributions, are poorly understood. Owen-Smith (1987) proposed the ‘keystone herbivore’ hypothesis, which provides a framework to explain the simultaneous extinctions of animals not obviously made extinct by the previous causes. Additionally, a possible cosmic impact may have generated the Younger Dryas cooling oscillation (12.8 and 11.0 k cal yr BP), resulting in a rapid defaunation process, including smaller taxa



Figure 3. Artistic reconstruction of *Onthophagus pilaus* nov. and its palaeoenvironment (by Mauricio Alvarez).

(Firestone et al. 2007; Pino et al. 2019; Wolbach et al. 2020). In this case, we suggest that both the rapid climatic changes and the extensive defaunaution in South America could be determining factors in the extinction of *O. pilaus* during the Pleistocene–Holocene transition.

Acknowledgements

We would like to thank to Dr Mario Pino for his collaboration in the manuscript revision, and Mario Elgueta for the primary approximation to the taxonomical identities. We are grateful to Fernanda Torres for the creation of the Figure 1B. FT was supported by CONICYT doctoral scholarship Grant #21171980 and Ilustre Municipalidad de Osorno, Osorno, Chile.

References

- Abarzúa AM, Martel-Cea A, Lobos V (2020) Vegetation–Climate–Megafauna Interactions During the Late Glacial in Pilauco Site, Northwestern Patagonia. In: Pino M, Astorga G (Eds) Pilauco: A Late Pleistocene Archaeo-paleontological Site. Springer, Cham, 157–173. https://doi.org/10.1007/978-3-030-23918-3_9
- Ashworth AC, Nelson RE (2014) The paleoenvironment of the Olympia beds based on fossil beetles from Discovery Park, Seattle, Washington, U.S.A. *Quaternary International* 341: 243–254 <https://doi.org/10.1016/j.quaint.2013.09.022>
- Ashworth AC, Hoganson J, Gunderson M (1989) Fossil-beetle analysis. In: Dillehay T (Ed.) Monte Verde – a Late Pleistocene Settlement in Chile. – a Late Pleistocene Settlement in Chile. Smithsonian Institution Press, Washington DC, 211–226.
- Barnosky AD, Koch PL, Feranec RS, Wing SL, Shabel AB (2004) Assessing the causes of late pleistocene extinctions on the continents. *Science* 306: 703–710 <https://doi.org/10.1126/science.1101476>
- Barnosky AD, Lindsey EL, Villavicencio NA, Bostelmann E, Hadly EA, Wanket J, Marshall CR (2016) Variable impact of late-Quaternary megafaunal extinction in causing ecological state shifts in North and South America. *Proceedings of the National Academy of Sciences* 113: 856–861 <https://doi.org/10.1073/pnas.1505295112>
- Borrero LA, Zárate M, Miotti L, Massone M (1998) The Pleistocene-Holocene transition and human occupations in the southern cone of South America. *Quaternary International* 49–50: 191–199 [https://doi.org/10.1016/S1040-6182\(97\)00063-3](https://doi.org/10.1016/S1040-6182(97)00063-3)
- Cantil LF, Sánchez MV, Bellosi ES, González MG, Sarzetti LC, Genise JC (2018) *Sphaera akatanaka* nov.: the first fossil brood ball attributable to necrophagous dung beetles associated with an Early Pleistocene environmental stress in the Pampean region (Argentina). *Palaeogeography, Palaeoclimatology, Palaeoecology* 386: 541–554. doi.org/10.1016/j.palaeo.2013.06.021
- Dillehay T (1989) Monte Verde. A Late Pleistocene Settlement in Chile. Smithsonian Institution Press, Washington DC, 1071 pp.
- Dillehay TD, Ocampo C, Saavedra J, Sawakuchi AO, Vega RM, Pino M, Collins MB, Scott Cummings L, Arregui I, Villagran XS, Hartmann GA, Mella M, González A, Dix G (2015) New Archaeological Evidence for an Early Human Presence at Monte Verde, Chile. Hart JP (Ed.) *PLoS ONE* 10(12): e0145471. [27 pp] <https://doi.org/10.1371/journal.pone.0141923>
- Elgueta M (2000) Estado actual del conocimiento de los coleópteros de Chile (Insecta: Coleoptera). *PRIBES-2000: Proyecto para Iberoamérica de Entomología Sistemática* 17: 145–154.
- Firestone RB, West A, Kennett JP, Becker L, Bunch TE, Revay ZS, Schultz PH, Belgia T, Kennett DJ, Erlandson JM, Dickenson OJ, Goodyear AC, Harris RS, Howard GA, Kloosterman JB, Lechler P, Mayewski PA, Montgomery J, Poreda R, Darrah T, Que Hee SS, Smitha AR, Stich A, Topping W, Wittke JH, Wolbach WS (2007) Evidence for an extraterrestrial impact 12,900 years ago that contributed to the megafaunal extinctions and the Younger Dryas cooling. *Proceedings of the National Academy of Sciences of the United States of America* 104: 16016–16021 <https://doi.org/10.1073/pnas.0706977104>
- González-Chang M, Pinochet D (2015) Escarabajos estercoleros nativos en Chile. Una revisión con énfasis en su ecología. *Agro Sur* 43: 511–521. <https://doi.org/10.4206/agrosur.2015.v43n3-06>

- González-Guarda E, Domingo L, Tornero C, Pino M, Hernández Fernández M, Sevilla P, Villavicencio N, Agustí J (2017) Late Pleistocene ecological, environmental and climatic reconstruction based on megafauna stable isotopes from northwestern Chilean Patagonia. *Quaternary Science Reviews* 170: 185–202. <https://doi.org/10.1016/j.quascirev.2017.06.035>
- Himes CMT, Gallardo MH, Kenagy GJ (2008) Historical biogeography and post-glacial recolonization of South American temperate rain forest by the relictual marsupial *Didelphis marsupialis*. *Journal of Biogeography* 35: 1415–1424. <https://doi.org/10.1111/j.1365-2699.2008.01895.x>
- Kuzmina SA, Korotyaev BA (2019) A new species of the weevil *Phyllotaxis* Germar, 1824 (Coleoptera: Curculionidae: Entiminae) from the Pleistocene of northeastern. *Invertebrate Zoology* 16: 154–164. <https://doi.org/10.15298/invertzool.16.2.04>
- Moreno K, Bostelmann JE, Macías C, Navarro-Harris X, De Pol-Holz R, Pino M (2019) A late Pleistocene human footprint from the Pilauco archaeological site, northern Patagonia, Chile. Chiang T-Y (Ed.) *PLoS ONE* 14(4): e0213572. <https://doi.org/10.1371/journal.pone.0213572>
- Navarro Harris X, Pino M, Guzmán Marín P, Lira MP, Labarca R, Corgne A (2019) Procurement and use of knappable glassy volcanic raw material from the late Pleistocene Pilauco site, Chilean Northwestern Patagonia. *Geoarchaeology* 34: 592–612. <https://doi.org/10.1002/gea.21736>
- Noriega J, Floate KD, Génier F, Reid CAM, Kohlmann B, Horgan FG, Davis ALV, Forgie SA, Aguilar C, Ibarra MG, Vaz de Mello F, Ziani S, Lobo JM (2020) Global distribution patterns provide evidence of niche shift by the introduced African dune *Dipodomys deserti* agus gazelle. *Entomologia Experimentalis et Applicata* 12961: 776–782. <https://doi.org/10.1111/eea.12961>
- Núñez JJ, Suárez-Villota EY, Quercia CA, Olivares AP, Sites Jr JW (2020) Phylogeographic analysis and species distribution modelling of the wood frog *Hyalinobatrachium* (Anura: Bufonidae) reveal interglacial diversification in south western Patagonia. *PeerJ* e9980: 1–29. <https://doi.org/10.7717/peerj.9980>
- Owen-Smith N (1987) Pleistocene Extinctions: the Pivotal Role of Megaherbivores. *Paleobiology* 13: 351–362. <https://doi.org/10.1017/S0094837300008927>
- Pino M, Abarzúa AM, Astorga G, Martel-Cea A, Cossio-Montecinos N, Navarro RX, Lira MP, Labarca R, LeCompte MA, Adedeji V, Moore CR, Bunch TE, Mooney C, Wolbach WS, West A, Kennett JP (2019) Sedimentary record from Patagonia, southern Chile supports cosmic-impact triggering of biomass burning, climate change, and megafaunal extinctions at 12.8 ka. *Scientific Reports* 9: 1–12. <https://doi.org/10.1038/s41598-018-38089-y>
- Pino M, Chávez-Homeister M, Navarro-Harris X, Labarca R (2013) The late Pleistocene Pilauco site, Osorno, south-central Chile. *Quaternary International* 299: 1–12. <https://doi.org/10.1016/j.quaint.2012.05.001>
- Pino M, Martel-Cea A, Vega RM, Fritte D, Soto-Bollmann K (2020) Geology, Stratigraphy, and Chronology of the Pilauco Site. In: Pino M, Astorga G (Eds) *Pilauco: A Late Pleistocene Archaeo-paleontological Site*. Springer, Cham, 303–312. https://doi.org/10.1007/978-3-030-23918-3_3

- Recabarren O (2020) e Proboscidean Gomphotheres (Mammalia, Gomphotheriidae) from Southernmost South America. In: Pino M, Astorga G (Eds) *Pilauco: A Late Pleistocene Archaeo-paleontological Site*. Springer, Cham, 546–548. https://doi.org/10.1007/978-3-030-23918-3_4
- Recabarren O, Pino M, Cid I (2011) A new record of *Equus* (Mammalia: Equidae) from the Late Pleistocene of central-south Chile. *Revista chilena de historia natural* 84: 535–542. <https://doi.org/10.4067/S0716-078X2011000400006>
- Rossini M, Vaz-de-Mello FZ, Zunino M (2018a) A taxonomic revision of the New World *Onthophagus* Latreille, 1802 (Coleoptera: Scarabaeidae: Scarabaeinae) species-complex, with description of two new species from South America. *Journal of Natural History* 52: 541–586. <https://doi.org/10.1080/00222933.2018.1437230>
- Rossini M, Vaz-De-Mello FZ, Zunino M (2018b) Toward a comprehensive taxonomic revision of the 'hirculus' group of american *Onthophagus* Latreille, 1802 (Coleoptera, Scarabaeidae, Scarabaeinae). *European Journal of Taxonomy* 432: 1–21. <https://doi.org/10.5852/ejt.2018.432>
- Sánchez MV, Genise JF, Bellosi ES, Román-Carrión JL, Cantil LF (2013) Dung beetle brood balls from Pleistocene highland palaeosols of Andean Ecuador: A reassessment of *Sauea* Coprinisphaera and their palaeoenvironments. *Palaeogeography, Palaeoclimatology, Palaeoecology* 386: 257–274. <https://doi.org/10.1016/j.palaeo.2013.05.028>
- Tarasov S, Dimitrov D (2016) Multigene phylogenetic analysis reveals dung beetles relationships and classification (Coleoptera: Scarabaeidae: Scarabaeinae). *BMC Evolutionary Biology* 16: 1–19. <https://doi.org/10.1186/s12862-016-0822-x>
- Tarasov S, Vaz-de-Mello FZ, Krell FT, Dimitrov D (2016) A review and phylogeny of Scarabaeine dung beetle fossils (Coleoptera: Scarabaeidae: Scarabaeinae), with the description of two *Canthochilum* species from Dominican amber. *PeerJ* 4:e1988. <https://doi.org/10.7717/peerj.1988>
- Tello F, Torres F (2020) Fossil Coleoptera from the Pilauco Site: An Approach to Late Pleistocene Microenvironments. In: Pino M, Astorga G (Eds) *Pilauco: A Late Pleistocene Archaeo-paleontological Site*. Springer, Switzerland, 195–212. https://doi.org/10.1007/978-3-030-23918-3_12
- Tello F, Arriagada G, Pino M (2019) First record of the family Histeridae (Insecta: Coleoptera) in a late Pleistocene sequence from Chile. *Ameghiniana* 57: 63–67. <https://doi.org/10.5710/AMGH.06.12.2019.3260>
- Tello F, Elgueta M, Abarzúa AM, Torres F, Pino M (2017) Fossil beetles from Pilauco, south-central Chile: An Upper Pleistocene paleoenvironmental reconstruction. *Quaternary International* 449: 58–66. <https://doi.org/10.1016/j.quaint.2017.05.046>
- Vaz-de-Mello FZ, Hal ter G (2006) A new dung beetle genus with two new species from Chile (Coleoptera: Scarabaeidae: Scarabaeinae). *Zootaxa* 1193: 1–6. <https://doi.org/10.11646/zootaxa.1193.1.4>
- Wolbach WS, Ballard JP, Mayewski PA, Kurbatov A, Bunch TE, LeCompte MA, Adedeji V, Israde-Alcántara I, Firestone RB, Mahaney WC, Melott AL, Moore CR, Napier WM, Howard GA, Tankersley KB, omas BC, Wittke JH, Johnson JR, Mitra S, Kennett JP, Kletetschka G, West A (2020) Extraordinary Biomass-Burning Episode and Impact Win

- ter Triggered by the Younger Dryas Cosmic Impact ~12,800 Years Ago: A Reply. *The Journal of Geology* 128: 95–107. <https://doi.org/10.1086/706265>
- Zinovyev E (2011) Sub-fossil beetle assemblages associated with the “mammoth fauna” in the Late Pleistocene localities of the Ural Mountains and West Siberia. *ZooKeys* 100: 149–169. <https://doi.org/10.3897/zookeys.100.1524>
- Zunino M (2013) Primer hallazgo de un escarabajo en un ichnofósil Coprinisphaeridae: *Phanaeus violetae* sp. (Coleoptera: Scarabaeinae) de una bola de cangahua del Ecuador. *Acta Zoológica Mexicana (N.S.)* 29: 219–226. <https://doi.org/10.21829/azm.2013.291397>
- Zunino M, Halter G (2019) About the origin of American Anthophagini (Coleoptera: Scarabaeidae). A critical appraisal of a recent paper by Breeschoten et al. (2016) on phylogenetics and biogeography. *Molecular Phylogenetics and Evolution* 133: 1411–1416. <https://doi.org/10.1016/j.ympev.2019.01.001>